

Morbidity/Mortality Topics	Indicator	Data Sources	Sub-group analysis	Timeframe	Benchmark	CSTE recommended surveillance indicators for substance abuse and mental health
Behavioral Health & substance abuse						
Morbidity	Suicide rate Liver Disease and cirrhosis	vital records vital records	Urban/rural, gender, age group (10-24, 25-64, 65+), race	10-year trend	US: HP 2020 MHMD-1 (LH)	Yes
	Drug overdose Opium, heroin, other opioids, methadone, other synthetic narcotics, cocaine, other and unspecified narcotics, cannabis, lysergide, other unspecified psychodjelentics, benzodiazepines, and psychostimulants Alcohol-related MMH crash deaths	vital records Fatality analysis reporting system (FARS)				Yes
	Hospitalizations (3 sub-indicators: Mood and depressive disorders, schizophrenic disorders, all mental disorders except drug- and alcohol-induced mental disorders)	MHDDS MHDDS (ED)	age group, urban/rural (or other geography), gender			Yes
	Prevalence: Substance Abuse Proportion of adolescents reporting use of alcohol or any illicit drugs during the past 30 days Proportion of persons engaging in binge drinking during the past 30 days (adults 18 years and older)	National Survey on Drug Use and Health (NSDUH); YRBS NSDUH; BRSS	Race, grade, gender Urban/Rural, race, gender, education, income		US: HP2020 SA-13.1 (LH) US: HP 2020 SA 14.3 (LH)	Yes Yes
	Dependence on or abuse of drugs or alcohol in the last year-- summary Use of selected prescription and illicit drugs (6 sub-indicators: illicit drug use in past month, marijuana use in past year, marijuana use in past month, illicit drug use other than marijuana use, 1 the past month, cocaine use in the past year, nonmedical use of pain relievers in past year)	NSDUH	Race, grade, gender			Yes
	Prevalence: Mental health Proportion of adolescents aged 12 to 17 years who experience major depressive episodes (MDEs)	National Survey on Drug Use and Health (NSDUH)			US: HP2020 MHMD-4.1 (LH)	Yes
	Prevalence of youth suicide attempts	YRBS	Race, grade, gender		US: HP2020 MHMD-2	Yes
	Prevalence of past-year serious mental illness	NSDUH	age group (218, 18-25, 26)		US: HP2020 MHMD-4.2	
	Prevalence of frequent mental distress (24 days out of 30)	NSDUH BRSS	age group (218, 18-25, 26) age group, sex, urban/rural, race			
Determinant/Correlater/Risk Factor Topics						
Topic	Indicator	Data Source	Sub-group analysis	Timeframe	Benchmark	Yes
Policy	State excise taxes on alcohol (for beer, wine, and spirits)	Alcohol Policy Information System (APIS)		2015		Yes
Sales	Prescription opioid sales per capita					Yes

*clear of information*